

Computing Solutions of Moral-Hazard Problems

Che-Lin Su

Dept. of Management Science and Engineering
Stanford University

Joint work with
Kenneth L. Judd
Hoover Institution
Stanford, CA 94305

July 19, 2005

Agenda

[Su & Judd 05] Computation of Moral-Hazard Problems

- Static Moral-Hazard Model
 - Deterministic contract
 - MPEC formulation
 - Hybrid method for a global solution
 - Numerical results on deterministic contract
 - Contract with action lotteries
 - Numerical results on Ned's example

Complementarity

Complementarity between nonnegative vectors: $x, y \in \mathbb{R}_+^n$

$$0 \leq x \perp y \geq 0 \iff x_i \geq 0, y_i \geq 0, x_i = 0 \text{ or } y_i = 0, \text{ for all } i$$

Complementarity

Complementarity between nonnegative vectors: $x, y \in \mathbb{R}_+^n$

$$0 \leq x \perp y \geq 0 \iff x_i \geq 0, y_i \geq 0, x_i = 0 \text{ or } y_i = 0, \text{ for all } i.$$

Complementarity Problem: Given a mapping $F : \mathbb{R}_+^n \rightarrow \mathbb{R}^n$,
find a vector $x \in \mathbb{R}_+^n$ satisfying

$$0 \leq x \perp F(x) \geq 0$$

Complementarity

Complementarity between nonnegative vectors: $x, y \in \mathbb{R}_+^n$

$$0 \leq x \perp y \geq 0 \iff x_i \geq 0, y_i \geq 0, x_i = 0 \text{ or } y_i = 0, \text{ for all } i.$$

Complementarity Problem: Given a mapping $F : \mathbb{R}_+^n \rightarrow \mathbb{R}^n$, find a vector $x \in \mathbb{R}_+^n$ satisfying

$$0 \leq x \perp F(x) \geq 0$$

Mathematical Program with Equilibrium Constraints (MPEC):

$$\begin{aligned} & \text{maximize}_{(x)} && f(x) \\ & \text{subject to} && g(x) \leq 0 \\ & && 0 \leq x \perp F(x) \geq 0 \end{aligned}$$

Difficulties with an MPEC


maximize $f(x, y)$

s.t. $x \geq 0$

$y \geq 0$


$xy = 0$

Difficulties with an MPEC




- **Nonconvex** feasible region.

Difficulties with an MPEC




- **Nonconvex** feasible region. Find a global solution?

Difficulties with an MPEC




- **Nonconvex** feasible region. Find a global solution?
- The standard CQs (LICQ, MFCQ) **fail** at every feasible point

Difficulties with an MPEC




- **Nonconvex** feasible region. Find a global solution?
- The standard CQs (LICQ, MFCQ) **fail** at every feasible point
 - Lagrangian multiplier set **unbounded**
 - Central path **does not exist**

Difficulties with an MPEC



- **Nonconvex** feasible region. Find a global solution?
- The standard CQs (LICQ, MFCQ) **fail** at every feasible point
 - Lagrangian multiplier set **unbounded**
 - Central path **does not exist**
- NLP solver **still work?**

Difficulties with an MPEC



- **Nonconvex** feasible region. Find a global solution?
- The standard CQs (LICQ, MFCQ) **fail** at every feasible point
 - Lagrangian multiplier set **unbounded**
 - Central path **does not exist**
- NLP solver **still work?** Yes, with modifications

Timeline of Moral-Hazard Model

- Notations:

- $a \in \mathcal{A} \subset \mathbb{R}$; $q_i \in \mathcal{Q} = \{q_1, \dots, q_N\}$; $c = (c_1, \dots, c_N) \in \mathbb{R}_+^N$

- Uncertainty: $p(q_i|a)$

- Utility: principal $w(q_i - c_i)$; agent $u(c_i, a)$

- Expected utility: principal $W(c, a) = \sum_i p(q_i|a)w(q_i - c_i)$;

agent $U(c, a) = \sum_i p(q_i|a)u(c_i, a)$

Timeline of Moral-Hazard Model

- Notations:

- $a \in \mathcal{A} \subset \mathbb{R}$; $q_i \in \mathcal{Q} = \{q_1, \dots, q_N\}$; $c = (c_1, \dots, c_N) \in \mathbb{R}_+^N$

- Uncertainty: $p(q_i|a)$

- Utility: principal $w(q_i - c_i)$; agent $u(c_i, a)$

- Expected utility: principal $W(c, a) = \sum_i p(q_i|a)w(q_i - c_i)$;

agent $U(c, a) = \sum_i p(q_i|a)u(c_i, a)$

- The principal and the agent agree to the contract with a suggested action a and compensation schedule $c = (c_1, \dots, c_N)$

Timeline of Moral-Hazard Model

- Notations:

- $a \in \mathcal{A} \subset \mathbb{R}$; $q_i \in \mathcal{Q} = \{q_1, \dots, q_N\}$; $c = (c_1, \dots, c_N) \in \mathbb{R}_+^N$

- Uncertainty: $p(q_i|a)$

- Utility: principal $w(q_i - c_i)$; agent $u(c_i, a)$

- Expected utility: principal $W(c, a) = \sum_i p(q_i|a)w(q_i - c_i)$;

agent $U(c, a) = \sum_i p(q_i|a)u(c_i, a)$

- The principal and the agent agree to the contract with a suggested action a and compensation schedule $c = (c_1, \dots, c_N)$
- The agent implements the suggested but unobservable action a

Timeline of Moral-Hazard Model

- Notations:

- $a \in \mathcal{A} \subset \mathbb{R}$; $q_i \in \mathcal{Q} = \{q_1, \dots, q_N\}$; $c = (c_1, \dots, c_N) \in \mathbb{R}_+^N$

- Uncertainty: $p(q_i|a)$

- Utility: principal $w(q_i - c_i)$; agent $u(c_i, a)$

- Expected utility: principal $W(c, a) = \sum_i p(q_i|a)w(q_i - c_i)$;

- agent $U(c, a) = \sum_i p(q_i|a)u(c_i, a)$

- The principal and the agent agree to the contract with a suggested action a and compensation schedule $c = (c_1, \dots, c_N)$
- The agent implements the suggested but unobservable action a
- Output q_i is realized

Timeline of Moral-Hazard Model

- Notations:

- $a \in \mathcal{A} \subset \mathbb{R}$; $q_i \in \mathcal{Q} = \{q_1, \dots, q_N\}$; $c = (c_1, \dots, c_N) \in \mathbb{R}_+^N$

- Uncertainty: $p(q_i|a)$

- Utility: principal $w(q_i - c_i)$; agent $u(c_i, a)$

- Expected utility: principal $W(c, a) = \sum_i p(q_i|a)w(q_i - c_i)$;

- agent $U(c, a) = \sum_i p(q_i|a)u(c_i, a)$

- The principal and the agent agree to the contract with a suggested action a and compensation schedule $c = (c_1, \dots, c_N)$
- The agent implements the suggested but unobservable action a
- Output q_i is realized
- Compensation c_i is paid according to the realized output q_i

Optimal Deterministic Contract

$$\begin{aligned} & \text{maximize}_{(c,a)} && W(c, a) \\ & \text{subject to} && U(c, a) \geq U_0 \text{ (Reservation Utility)} \\ & && a \in \operatorname{argmax}\{U(c, \tilde{a}) : \tilde{a} \in \mathcal{A}\} \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

Optimal Deterministic Contract

$$\begin{aligned} & \text{maximize}_{(c,a)} && W(c, a) \\ & \text{subject to} && U(c, a) \geq U_0 \text{ (Reservation Utility)} \\ & && a \in \operatorname{argmax}\{U(c, \tilde{a}) : \tilde{a} \in \mathcal{A}\} \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

- Both c and a are continuous variables
- Need global optimality at both levels

Optimal Deterministic Contract

$$\begin{aligned} & \text{maximize}_{(c,a)} && W(c, a) \\ & \text{subject to} && U(c, a) \geq U_0 \text{ (Reservation Utility)} \\ & && a \in \operatorname{argmax}\{U(c, \tilde{a}) : \tilde{a} \in \mathcal{A}\} \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

- Both c and a are **continuous** variables
- Need **global optimality** at **both** levels
- First-order approach [Mirrlees '75] [Rogerson '85]
 - Replace $\{\max U(c, a) : a \in \mathcal{A}\}$ by first-order conditions, but usually, $U(c, a)$ is **not** concave in a for all c

LP Lotteries Approach

- [Myerson '79] [Prescott & Townsend '84a, b] [(Ned) Prescott '04]
- Consider **finite action** set \mathcal{A} and **finite compensation** set \mathcal{C} with element ξ

LP Lotteries Approach

- [Myerson '79] [Prescott & Townsend '84a, b] [(Ned) Prescott '04]
- Consider **finite action** set \mathcal{A} and **finite compensation** set \mathcal{C} with element ξ
- Timeline
 - The principal and the agent agree to the contract with lotteries $\pi(a)$ and $\pi(\xi|q, a)$

LP Lotteries Approach

- [Myerson '79] [Prescott & Townsend '84a, b] [(Ned) Prescott '04]
- Consider **finite action** set \mathcal{A} and **finite compensation** set \mathcal{C} with element ξ
- Timeline
 - The principal and the agent agree to the contract with lotteries $\pi(a)$ and $\pi(\xi|q, a)$
 - The action lottery is done according to the agreed probabilities $\pi(a)$

LP Lotteries Approach

- [Myerson '79] [Prescott & Townsend '84a, b] [(Ned) Prescott '04]
- Consider **finite action** set \mathcal{A} and **finite compensation** set \mathcal{C} with element ξ
- Timeline
 - The principal and the agent agree to the contract with lotteries $\pi(a)$ and $\pi(\xi|q, a)$
 - The action lottery is done according to the agreed probabilities $\pi(a)$
 - Output q is realized

LP Lotteries Approach

- [Myerson '79] [Prescott & Townsend '84a, b] [(Ned) Prescott '04]
- Consider **finite action** set \mathcal{A} and **finite compensation** set \mathcal{C} with element ξ
- Timeline
 - The principal and the agent agree to the contract with lotteries $\pi(a)$ and $\pi(\xi|q, a)$
 - The action lottery is done according to the agreed probabilities $\pi(a)$
 - Output q is realized
 - Compensation is paid according to the agreed lottery conditional on the action a and the realized output q : $\pi(\xi|q, a)$

LP Lotteries Approach

- Ideas: Consider **action** and **compensation** lotteries
 - Construct an action grid $\mathcal{A} = \{a_1, \dots, a_M\}$ and a compensation grid $\mathcal{C} = \{\xi_1, \dots, \xi_L\}$
 - Introduce probability measures $\pi(a)$ and $\pi(\xi|q, a)$
 - Transform ODC into an **LP** with $\pi = (\pi(\xi, q, a))_{\xi \in \mathcal{C}, q \in \mathcal{Q}, a \in \mathcal{A}}$

LP Lotteries Approach

- Ideas: Consider **action** and **compensation** lotteries
 - Construct an action grid $\mathcal{A} = \{a_1, \dots, a_M\}$ and a compensation grid $\mathcal{C} = \{\xi_1, \dots, \xi_L\}$
 - Introduce probability measures $\pi(a)$ and $\pi(\xi|q, a)$
 - Transform ODC into an **LP** with $\pi = (\pi(\xi, q, a))_{\xi \in \mathcal{C}, q \in \mathcal{Q}, a \in \mathcal{A}}$
- Disadvantages:
 - The resulting LP is **large**:
 $L * M * N$ variables and $M * (N + M - 1) + 2$ constraints.
If $M = 50, N = 40, L = 500, \Rightarrow$ an LP with **one million** variables and **4452** constraints
 - Economic choice variables are continuous in nature
 - The curse of dimensionality for multidimensional problems

MPEC Approach for finite \mathcal{A}

- Goal of [Su-Judd '05]: Allow continuous compensation $c \in \mathbb{R}_+^N$ (and keep finite \mathcal{A})

MPEC Approach for finite \mathcal{A}

- Goal of [Su-Judd '05]: Allow continuous compensation $c \in \mathbb{R}_+^N$ (and keep finite \mathcal{A})
- Idea: Introduce mixed-strategy profile $\delta = (\delta_1, \dots, \delta_M)$ for agent's action choices $\mathcal{A} = \{a_1, \dots, a_M\}$:

$$\delta_k \iff a_k \quad \text{for } k = 1, \dots, M$$

MPEC Approach for finite \mathcal{A}

- Goal of [Su-Judd '05]: Allow continuous compensation $c \in \mathbb{R}_+^N$ (and keep finite \mathcal{A})
- Idea: Introduce mixed-strategy profile $\delta = (\delta_1, \dots, \delta_M)$ for agent's action choices $\mathcal{A} = \{a_1, \dots, a_M\}$:

$$\delta_k \iff a_k \quad \text{for } k = 1, \dots, M$$

- Agent's mixed strategy problem is an LP

$$a^* \in \operatorname{argmax} \{U(c, a) : a \in \{a_1, \dots, a_M\}\}$$
$$\Downarrow$$
$$\delta^* \in \operatorname{argmax} \left\{ \sum_{k=1}^M U(c, a_k) \delta_k : e^T \delta = 1, \delta \geq 0 \right\}$$

MPEC Approach for finite \mathcal{A}

- Goal of [Su-Judd '05]: Allow continuous compensation $c \in \mathbb{R}_+^N$ (and keep finite \mathcal{A})
- Idea: Introduce mixed-strategy profile $\delta = (\delta_1, \dots, \delta_M)$ for agent's action choices $\mathcal{A} = \{a_1, \dots, a_M\}$:

$$\delta_k \iff a_k \quad \text{for } k = 1, \dots, M$$

- Agent's mixed strategy problem is an LP

$$a^* \in \operatorname{argmax} \{U(c, a) : a \in \{a_1, \dots, a_M\}\}$$

$$\delta^* \in \operatorname{argmax} \left\{ \sum_{k=1}^M U(c, a_k) \delta_k : e^T \delta = 1, \delta \geq 0 \right\}$$

Optimality + strong duality \iff Let $U(c) := (U(c, a_k))_{k=1}^M$

$$\delta^* \text{ solves } \begin{cases} 0 \leq \delta \perp (U(c)^T \delta) e - U(c) \geq 0 \\ e^T \delta = 1 \end{cases}$$

MPEC Approach for finite \mathcal{A}

$$\text{maximize}_{(c,a)} \quad W(c, a)$$

$$\text{subject to} \quad U(c, a) \geq U_0$$

$$a \in \operatorname{argmax} \{U(c, \tilde{a}) : \tilde{a} \in \mathcal{A} = \{a_1, \dots, a_M\}\}$$

$$c \in \mathbb{R}_+^N$$

MPEC Approach for finite \mathcal{A}

$$\begin{aligned} & \text{maximize}_{(c, \delta)} && W(c)^T \delta \\ & \text{subject to} && U(c)^T \delta \geq U_0 \\ & && e^T \delta = 1 \\ & && 0 \leq \delta \perp (U(c)^T \delta) e - U(c) \geq 0 \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

MPEC Approach for finite \mathcal{A}

$$\begin{aligned} & \text{maximize}_{(c, \delta)} && W(c)^T \delta \\ & \text{subject to} && U(c)^T \delta \geq U_0 \\ & && e^T \delta = 1 \\ & && 0 \leq \delta \perp (U(c)^T \delta) \quad e - U(c) \geq 0 \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

This is a **M**athematical **P**rogram with **E**quilibrium **C**onstraints (**MPEC**)!

MPEC Approach for finite \mathcal{A}

$$\begin{aligned} & \text{maximize}_{(c, \delta)} && W(c)^T \delta \\ & \text{subject to} && U(c)^T \delta \geq U_0 \\ & && e^T \delta = 1 \\ & && 0 \leq \delta \perp (U(c)^T \delta) e - U(c) \geq 0 \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

This is a **M**athematical **P**rogram with **E**quilibrium **C**onstraints (**MPEC**)!

Problem Size: $(N + M)$ variables, $2M + 2$ constraints

MPEC Approach for finite \mathcal{A}

$$\begin{aligned} & \text{maximize}_{(c, \delta)} && W(c)^T \delta \\ & \text{subject to} && U(c)^T \delta \geq U_0 \\ & && e^T \delta = 1 \\ & && 0 \leq \delta \perp (U(c)^T \delta) e - U(c) \geq 0 \\ & && c \in \mathbb{R}_+^N \end{aligned}$$

This is a **M**athematical **P**rogram with **E**quilibrium **C**onstraints (**MPEC**)!

Problem Size: $(N + M)$ variables, $2M + 2$ constraints

Lemma:

- (i) $(c^*, \delta^*) \in \mathbf{SOL}(\text{MPEC}) \implies (c^*, a_i^*)$ an ODC with $a_i^* \in \mathcal{A}$ finite, and $i \in \{j : \delta_j^* > 0\}$.
- (ii) (c^*, a_i^*) an ODC with $a_i^* \in \mathcal{A}$ finite $\implies (c^*, e_i) \in \mathbf{SOL}(\text{MPEC})$, where e_i is the i -th column of an identity matrix

A Hybrid Method [Su-Judd '05]

Observations:

- MPEC: Allows continuous compensation $c \in \mathbb{R}_+^N$ but may stop at local maximum
- LP: Produce global optimal but the grid \mathcal{C} may be too coarse

A Hybrid Method [Su-Judd '05]

Observations:

- MPEC: Allows continuous compensation $c \in \mathbb{R}_+^N$ but may stop at local maximum
- LP: Produce global optimal but the grid \mathcal{C} may be too coarse

Goal: Combine the best features of both to find a **global** solution with continuous compensation $c \in \mathbb{R}_+^N$

A Hybrid Method [Su-Judd '05]

Observations:

- MPEC: Allows continuous compensation $c \in \mathbb{R}_+^N$ but may stop at local maximum
- LP: Produce global optimal but the grid \mathcal{C} may be too coarse

Goal: Combine the best features of both to find a global solution with continuous compensation $c \in \mathbb{R}_+^N$

Step 0: Construct a coarse grid \mathcal{C}

Step 1: Solve the **LP** for the given grid \mathcal{C}

Step 2: Setup and solve the **MPEC** :

$$\begin{cases} (2.1) : & \text{construct } (c^0, \delta^0) \text{ using LP solution} \\ (2.2) : & \text{Solve the MPEC with the starting point } (c^0, \delta^0) \end{cases}$$

Step 3: Refine the grid and repeat Step 1 and Step 2

A Hybrid Method [Su-Judd '05]

Observations:

- MPEC: Allows continuous compensation $c \in \mathbb{R}_+^N$ but may stop at local maximum
- LP: Produce global optimal but the grid \mathcal{C} may be too coarse

Goal: Combine the best features of both to find a global solution with continuous compensation $c \in \mathbb{R}_+^N$

Step 0: Construct a coarse grid \mathcal{C}

Step 1: Solve the **LP** for the given grid \mathcal{C}

Step 2: Setup and solve the **MPEC** :

$$\begin{cases} (2.1) : & \text{construct } (c^0, \delta^0) \text{ using LP solution} \\ (2.2) : & \text{Solve the MPEC with the starting point } (c^0, \delta^0) \end{cases}$$

Step 3: Refine the grid and repeat Step 1 and Step 2

Result: A hybrid solution is always better than an LP solution

An Example in [Karaivanov '01]

- Risk-neutral principal: $w(q_i - c_i) = q_i - c_i$
- Risk-averse agent: $u(c_i, a) = \frac{c_i^{1-\gamma}}{1-\gamma} + \kappa \frac{(1-a)^{1-\delta}}{1-\delta}$
- Two outcomes: $q_H = \$3$ and $q_L = \$1$
- Action set: $|\mathcal{A}| = 10$ with equally-spaced effort level within $[0.01, 0.99]$
- The production technology $p(q = q_H|a) = a^\alpha$ with $0 < \alpha < 1$

γ	κ	δ	α	U_0
0.5	1	0.5	0.7	1

- Both LP and MPEC are coded in AMPL and solved by SNOPT on NEOS server (host: `prado.iems.northwestern.edu`)

LP Solutions

LP Solutions with 8 different grids (# of constraints = 112)

$ C $	# of Variables	Read Time (in sec.)	Solve Time (in sec.)	# of Iterations	Objective Value
21	420	0.01	0.04	37	1.875882746
41	820	0.02	0.07	46	1.877252910
81	1620	0.03	0.12	46	1.877259193
161	3220	0.06	0.25	46	1.877262265
321	6420	0.13	0.58	69	1.877263785
641	12820	0.26	1.12	52	1.877259905
1281	25620	0.53	2.67	101	1.877262221
2561	51220	1.09	4.81	73	1.877262201
5121	102420	2.46	11.70	101	1.877263113

Hybrid Solution

LP $ C $	Read Time (in sec.)	Solve Time (in sec.)	# of Iterations	Objective Value
21	0.01	0.04	37	1.875882746
MPEC Starting Point	Read Time (in sec.)	Solve Time (in sec.)	# of Major Iterations	Objective Value
$\delta_6 = 1, \delta_{i(\neq 6)} = 0$	0.02	0.01	13	1.877265298

Hybrid Solution

LP $ \mathcal{C} $	Read Time (in sec.)	Solve Time (in sec.)	# of Iterations	Objective Value
21	0.01	0.04	37	1.875882746
MPEC Starting Point	Read Time (in sec.)	Solve Time (in sec.)	# of Major Iterations	Objective Value
$\delta_6 = 1, \delta_{i(\neq 6)} = 0$	0.02	0.01	13	1.877265298

LP solution

LP $ \mathcal{C} $	Read Time (in sec.)	Solve Time (in sec.)	# of Iterations	Objective Value
5121	2.46	11.70	101	1.877263113

Contract with Action Lotteries

- A probability distribution $\pi(a)$ and a compensation schedule $c(a) \in R^N$ for every action $a \in \mathcal{A}$

Contract with Action Lotteries

- A probability distribution $\pi(a)$ and a compensation schedule $c(a) \in R^N$ for every action $a \in \mathcal{A}$
- Modifications:
 - Objective function: $\sum_a \pi(a)W(c(a), a)$
 - Participation constraint: $\sum_a \pi(a)U(c(a), a) \geq U_0$
 - Incentive compatibility constraints:

$$\text{if } \pi(a) > 0, \text{ then } \begin{cases} \pi(a) \geq 0 \\ U(c(a), a) \geq U(c(a), a_1), \\ \vdots \\ U(c(a), a) \geq U(c(a), a_M). \end{cases}$$

$$\Updownarrow$$

$$\pi(a) \geq 0$$

$$\pi(a)(U(c(a), a) - U(c(a), \tilde{a})) \geq 0, \quad \forall (a, \tilde{a}) \in \mathcal{A} \times \mathcal{A}$$

Contract with Action Lotteries

- A probability distribution $\pi(a)$ and a compensation schedule $c(a) \in R^N$ for every action $a \in \mathcal{A}$

$$\text{maximize } \sum_{a \in \mathcal{A}} \pi(a) W(c(a), a)$$


$$\text{subject to } \sum_{a \in \mathcal{A}} \pi(a) U(c(a), a) \geq U_0,$$

$$\sum_{a \in \mathcal{A}} \pi(a) = 1,$$


$$\pi(a) (U(c(a), a) - U(c(a), \tilde{a})) \geq 0, \quad \forall (a, \tilde{a} (\neq a)) \in \mathcal{A} \times \mathcal{A},$$

$$\pi(a) \geq 0, \quad \forall a \in \mathcal{A}.$$

Star-Shaped Feasible Region



Star-Shaped Feasible Region



- **Nonconvex** feasible region [Scholtes '04]
- LICQ **fails** on y -axis

MPEC for Action Lotteries

$$\pi(a) \geq 0, \quad \text{if } \pi(a) > 0, \text{ then } \begin{cases} U(c(a), a) \geq U(c(a), a_1), \\ \vdots \\ U(c(a), a) \geq U(c(a), a_M). \end{cases}$$

\Leftrightarrow

$$\forall a \in \mathcal{A}: \begin{cases} \pi(a) (U(c(a), a) - U(c(a), \tilde{a})) \geq 0, & \forall \tilde{a} (\neq a) \in \mathcal{A}, \\ \pi(a) \geq 0. \end{cases}$$

\Leftrightarrow

$$\forall a \in \mathcal{A}: \begin{cases} U(c(a), a) - U(c(a), \tilde{a}) + s(a) \geq 0, & \forall \tilde{a} (\neq a) \in \mathcal{A}, \\ 0 \leq \pi(a) \perp s(a) \geq 0. \end{cases}$$

MPEC for Action Lotteries

$$\text{maximize } \sum_{a \in \mathcal{A}} \pi(a) W(c(a), a)$$

$$\text{subject to } \sum_{a \in \mathcal{A}} \pi(a) U(c(a), a) \geq U_0,$$

$$\sum_{a \in \mathcal{A}} \pi(a) = 1,$$

$$\forall a \in \mathcal{A} : \begin{cases} U(c(a), a) - U(c(a), \tilde{a}) + s(a) \geq 0, & \forall \tilde{a} (\neq a) \in \mathcal{A}, \\ 0 \leq \pi(a) \perp s(a) \geq 0. \end{cases}$$

The A-L Example in [Prescott '04]

Hybrid method: LP with 11 compensation grid points and then switch to MPEC.

Hybrid Solutions with 10 different action grids, $|\mathcal{C}| = 11$, $|\mathcal{Q}| = 50$

$ \mathcal{A} $	t_{LP} (in sec.)	LP Obj. Val.	t_{MPEC} (in sec.)	MPEC Obj. Val.	t_{Total} (in sec.)
6	2	1.75868508	1	1.76234445	3
11	4	1.75868508	2	1.76234448	6
16	9	1.76265085	3	1.76565622	12
21	12	1.76351860	11	1.76630661	23
26	21	1.75924445	6	1.76298273	27
31	38	1.76265085	11	1.76565620	49
36	64	1.75606325	16	1.76051776	80
51	102	1.75924445	22	1.76298271	124
76	266	1.75778364	127	1.76298273	393
101	575	1.75572408	1108	1.76234445	1683
151	4203	1.75534000	14018	1.75996707	18221

- Plan to use penalty approach to solve large MPECs (see Sven's talk).

Extensions

- MPEC formulations are also given for :
 - Contracts with compensation lottery (randomized payment)
 - Contracts with action and compensation lotteries
 - Multidimensional action choices - little economic theory, make special assumptions
 - Multidimensional compensation choices - infeasible for LP

Extensions

- MPEC formulations are also given for :
 - Contracts with compensation lottery (randomized payment)
 - Contracts with action and compensation lotteries
 - Multidimensional action choices - little economic theory, make special assumptions
 - Multidimensional compensation choices - infeasible for LP
- Future research
 - Tournament (single-principal multi-agent) problem [Lazear & Rosen '81]
 - Incentive problem with both adverse selection and moral-hazard
 - Dynamic contracts (multi-period moral-hazard problem) [Phelan & Townsend '91]
 - Star-shaped problems [Scholtes '04]

Optimal Income Taxation with Multidimensional Types

Kenneth L. Judd

Che-Lin Su

Hoover Institution

Stanford University

PRELIMINARY AND INCOMPLETE

July, 2005

Introduction

- Optimal income taxation
 - Mirrlees-style analysis
 - * Heterogeneous productivity
 - * Common utility function
 - * Utilitarian (or redistributive) objective
 - * Optimal tax rates are high initially but fall; asymptotically zero for thin tailed distributions of ability
 - * Clear pattern of binding IC constraints
 - Example of nonlinear optimization with incentive compatibility constraints
- Multidimensional heterogeneity
 - Little theory; special cases only
 - No clear pattern of binding IC constraints
 - Revelation principle still holds, producing a nonlinear optimization problem with IC constraints.
 - Clearly more realistic than 1-D models.

- This paper examines two-dimensional heterogeneity
 - Heterogeneous ability and elasticity of labor supply
 - We take a numerical approach
 - * This is not as difficult as commonly perceived.
 - * Use various diagnostics to avoid spurious local minima - result is more reliable than standard ML estimation.
 - Optimal marginal tax rate at top can be significantly negative
 - Binding incentive constraints are **not** local.

Model

- N types of taxpayers.
- Two goods: consumption and labour services. Let c_i (l_i) denote taxpayer i 's consumption (labor supply).
- Taxpayer i 's productivity is w_i ; $0 < w_1 < \dots < w_N$, i 's pretax income is

$$y_i := w_i l_i, \quad i = 1, \dots, N \quad (1)$$

- Define utility function $U^i : R \times R_+ \rightarrow R$ over income and consumption - the observables

$$U^i(y_i, c_i) := u_i(c_i, y_i/w_i), \quad i = 1, \dots, N. \quad (2)$$

- The *social welfare function* $W : R^N \times R_+^N, \rightarrow R$ is

$$W(a) := \sum_i \lambda_i U^i(y_i, c_i), \quad (3)$$

where λ_i equals the population frequency of type i : the utilitarian social welfare function. Also, $\sum_i \lambda_i = 1$.

- Resource constraint: $\sum_i c_i \leq \sum_i y_i$
- Anonymous taxation: each taxpayer can choose any (y_i, c_i) bundle offered by the government.
- Revelation principle: government constructs a schedule such that type i taxpayers will choose the (y_i, c_i) bundle
- Government problem is a nonlinear optimization problem with nonlinear constraints:

$$\begin{aligned} \max_{y_i, c_i} \quad & \sum_i \lambda_i u_i(c_i, y_i/w_i) \\ u_i(c_i, y_i/w_i) \geq & u_i(c_j, y_j/w_i), \forall i, j \\ \sum_i c_i \leq & \sum_i w_i l_i \\ \sum_i c_i \geq & 0 \end{aligned} \quad (4)$$

- The zero tax commodity bundles are the solutions to

$$\max_l u_i(w_i l, l)$$

for each type i . We denote the zero tax commodity bundles by (c^*, l^*, y^*) .

- Example:

$$u(c, l) = \log c - l^{\eta+1}/(\eta + 1)$$

$$N = 5, \lambda_i = 1/N$$

$$w_i \in \{1, 2, 3, 4, 5\}$$

- The zero tax solution is $l_i = 1, c_i = w_i$
- We compute the solutions for various w and η , and report the following:

$$ATR_i = \frac{y_i - c_i}{y_i}, \quad i = 1, \dots, N$$

$$MTR_i = 1 - \frac{u_l}{w u_c}, \quad i = 1, \dots, N - 1$$

Mirrlees Cases: common utility function

Table 1. $\eta = 1$						
i	c_i	y_i	$\frac{y_i - c_i}{y_i}$	MTR_i	l_i/l_i^*	c_i/c_i^*
1	1.56	0.40	-2.87	0.63	0.40	1.56
2	1.90	1.31	-0.45	0.53	0.65	0.95
3	2.49	2.56	0.03	0.40	0.85	0.83
4	3.36	4.01	0.16	0.25	1.00	0.84
5	4.50	5.54	0.19	–	1.10	0.90

Table 2. $\eta = 2$						
i	c_i	y_i	$\frac{y_i - c_i}{y_i}$	MTR_i	l_i/l_i^*	c_i/c_i^*
1	1.87	0.60	-2.09	0.68	0.60	1.87
2	2.16	1.54	-0.39	0.59	0.77	1.08
3	2.63	2.69	0.02	0.47	0.89	0.87
4	3.31	3.99	0.17	0.32	0.99	0.82
5	4.26	5.41	0.21	–	1.08	0.85

Table 3. $\eta = 3$:						
i	c_i	y_i	$\frac{y_i - c_i}{y_i}$	MTR_i	l_i/l_i^*	c_i/c_i^*
1	2.06	0.70	-1.91	0.73	0.70	2.06
2	2.31	1.66	-0.38	0.64	0.83	1.15
3	2.71	2.77	0.02	0.53	0.92	0.90
4	3.28	3.99	0.17	0.38	0.99	0.82
5	4.10	5.33	0.23	–	1.06	0.82

Table 4. $\eta = 5$						
i	c_i	y_i	$\frac{y_i - c_i}{y_i}$	MTR_i	l_i/l_i^*	c_i/c_i^*
1	2.29	0.80	-1.84	0.79	0.80	2.29
2	2.49	1.78	-0.39	0.71	0.89	1.24
3	2.80	2.85	0.02	0.61	0.95	0.93
4	3.24	4.01	0.19	0.48	1.00	0.81
5	3.88	5.25	0.26	–	1.05	0.77

Table 5. $\eta = 8$						
i	c_i	y_i	$\frac{y_i - c_i}{y_i}$	MTR_i	l_i/l_i^*	c_i/c_i^*
1	2.48	0.87	-1.84	0.84	0.87	2.48
2	2.63	1.86	-0.41	0.77	0.93	1.31
3	2.86	2.91	0.02	0.69	0.97	0.95
4	3.20	4.02	0.20	0.58	1.00	0.80
5	3.68	5.19	0.28	–	1.03	0.73

Two-Dimensional Types - Productivity and Elasticity of Labor Supply

- $u^j(c, l) = \log c - l^{\eta_j+1}/(\eta_j + 1)$
- w_i is productivity type i .
- (c_{ij}, y_{ij}) is allocation for (i, j) -type taxpayer.
- Zero tax solution for type (i, j) is $(l_{ij}^*, c_{ij}^*, y_{ij}^*) = (1, w_i, w_i)$.
- Problem:

$$\begin{aligned} \max_{(y,c)} \quad & \sum_{i=1}^N \sum_{j=1}^N \lambda_{ij} u^j(c_{ij}, y_{ij}/w_i) \\ & u^j(c_{ij}, y_{ij}/w_i) - u^j(c_{i'j'}, y_{i'j'}/w_i) \geq 0 \quad \forall (i, j), (i', j') \\ & \sum_{i=1}^N \sum_{j=1}^N c_{ij} \leq \sum_{i=1}^N y_{ij} \\ & \sum_{i=1}^N \sum_{j=1}^N c_{ij} \geq 0, \end{aligned}$$

- We choose the following parameters:
 - $N = 5, w_i = i$
 - $\lambda_i = 1/(N^2)$.
 - $\eta = (1, 2, 3, 5, 8)$.
 - We use the zero tax solution (c^*, y^*) as a starting point for the NLP solver.

Table 6.: $\eta = (1, 2, 3, 5, 8)$, $w = (1, 2, 3, 4, 5)$

(i, j)	c_{ij}	y_{ij}	$\Delta TR_{i,j}$	$MTR_{i,j}$	$ATR_{i,j}$	l_{ij}/l_{ij}^*	c_{ij}/c_{ij}^*	Utility	
								Judd-Su	Mirrlees
(1, 1)	1.68	0.42		0.28	-2.92	0.42	1.68	0.4294	.3641
(1, 2)	1.77	0.62	0.51	0.32	-1.86	0.62	1.77	0.4952	.3138
(1, 3)	1.79	0.65	0.54	0.51	-1.75	0.65	1.79	0.5378	.6601
(1, 4)	1.83	0.77	0.66	0.50	-1.37	0.77	1.83	0.5700	.7830
(1, 5)	1.86	0.86	0.62	0.43	-1.16	0.86	1.86	0.5940	.8760
(2, 1)	1.86	0.86		0.60	-1.16	0.43	0.93	0.5308	.3751
(2, 2)	2.03	1.39	0.68	0.50	-0.45	0.69	1.01	0.5973	.6180
(2, 3)	2.07	1.50	0.60	0.56	-0.38	0.75	1.03	0.6512	.7189
(2, 4)	2.16	1.74	0.62	0.46	-0.24	0.87	1.08	0.7006	.8181
(2, 5)	2.20	1.83	0.55	0.46	-0.20	0.91	1.10	0.7413	.9085
(3, 1)	2.20	1.83		0.55	-0.20	0.61	0.73	0.6053	.5496
(3, 2)	2.47	2.49	0.59	0.43	0.00	0.83	0.82	0.7157	.7269
(3, 3)	2.47	2.49		0.53	0.00	0.83	0.82	0.7878	.8158
(3, 4)	2.55	2.68	0.59	0.52	0.04	0.89	0.85	0.8520	.9057
(3, 5)	2.62	2.85	0.54	0.42	0.07	0.95	0.87	0.8965	.9672
(4, 1)	3.36	4.00	0.36	0.16	0.15	1.00	0.84	0.7127	.7090
(4, 2)	3.36	4.00	–	0.16	0.15	1.00	0.84	0.8794	.8664
(4, 3)	3.36	4.00	–	0.15	0.15	1.00	0.84	0.9627	.9402
(4, 4)	3.36	4.00	–	0.15	0.15	1.00	0.84	1.0461	1.0080
(4, 5)	3.36	4.00		0.15	0.15	1.00	0.84	1.1017	1.0476
(5, 5)	4.00	5.14	0.44	0	0.22	1.02	0.80	1.2439	1.1487
(5, 4)	4.11	5.24	-0.10	-0.05	0.21	1.04	0.82	1.1928	1.1331
(5, 3)	4.34	5.43	-0.17	-0.12	0.20	1.08	0.86	1.1188	1.0877
(5, 2)	4.49	5.56	-0.17	-0.11	0.19	1.11	0.89	1.0428	1.0286
(5, 1)	4.87	5.87	-0.22	-0.15	0.17	1.17	0.97	0.8933	.8901

Table 7. Binding IC $[(i, j), (i', j')]$

(i, j)	(i', j')	(i, j)	(i', j')
(1, 2)	(1, 1)	(4, 1)	(3, 2), (3, 3), (3, 5), (4, 2), (4, 3), (4, 4), (4, 5)
(1, 3)	(1, 2)	(4, 2)	(4, 1), (4, 3), (4, 4), (4, 5)
(1, 4)	(1, 3)	(4, 3)	(4, 1), (4, 2), (4, 4), (4, 5)
(1, 5)	(1, 4), (2, 1)	(4, 4)	(4, 1), (4, 2), (4, 3), (4, 5)
(2, 1)	(1, 4), (1, 5)	(4, 5)	(4, 1), (4, 2), (4, 3), (4, 4)
(2, 2)	(1, 5), (2, 1)	(5, 1)	(4, 1), (4, 2), (4, 3), (4, 4), (4, 5)
(2, 3)	(2, 2)	(5, 2)	(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (5, 1)
(2, 4)	(2, 3)	(5, 3)	(5, 2)
(2, 5)	(2, 4), (3, 1)	(5, 4)	(5, 3)
(3, 1)	(2, 3), (2, 5)	(5, 5)	(5, 4)
(3, 2)	(2, 5), (3, 1), (3, 3)		
(3, 3)	(3, 2)		
(3, 4)	(3, 2), (3, 3)		
(3, 5)	(3, 4)		

Comparisons

- Negative marginal rates at top in heterogeneous η . case
- Binding IC constraints are not local in income
- Less redistribution in heterogeneous η case
 - Average tax rates are lower for top two productivity types
 - Marginal tax rates are lower for top two productivity types
- More output - both consumption and labor supply tends to be higher

Robustness

- Rawlsian objective
 - Negative marginal rates at top
 - Rich do not do much worse, if at all
- Government expenditure
 - Efficiency becomes more important
 - Less redistribution
 - Negative rates at top persist
- Log normal distributed abilities
 - More realistic
 - Same patterns of results
- Number of types – 15 w 's and 5 η 's give same results

Two-Dimensional Types - Productivity and Labor Disutility

- $u^j(c, l) = \log c - \eta_j l^2/2$
- w_i is productivity type i .
- (c_{ij}, y_{ij}) is allocation for (i, j) -type taxpayer.
- Problem:

$$\begin{aligned} \max_{(y,c)} \quad & \sum_{i=1}^N \sum_{j=1}^N \lambda_{ij} u^j(c_{ij}, y_{ij}/w_i) \\ & u^j(c_{ij}, y_{ij}/w_i) - u^j(c_{i'j'}, y_{i'j'}/w_i) \geq 0 \quad \forall (i, j), (i', j') \\ & \sum_{i=1}^N \sum_{j=1}^N c_{ij} \leq \sum_{i=1}^N y_{ij} \\ & \sum_{i=1}^N \sum_{j=1}^N c_{ij} \geq 0, \end{aligned}$$

- We choose the following parameters:
 - $N = 5$, $w_i = i$
 - $\eta = (1/8, 1/4, 1/2, 3/4, 1)$.
 - We use the zero tax solution (c^*, y^*) as a starting point for the NLP solver.

Table 8: Heterogeneous wages and tastes for leisure

i	η	c	y	ATR	MTR	l_{ij}/l_{ij}^*	c_{ij}/c_{ij}^*
1	1	1.591	0.256	-5.20	0.18	0.36	2.25
1	3/4	1.595	0.261	-5.09	0.37	0.32	1.95
1	1/2	1.597	0.267	-4.97	0.57	0.26	1.59
1	1/4	1.773	0.699	-1.53	0.37	0.49	1.25
2	3/4	1.773	0.699	-1.53	0.53	0.42	1.08
2	1	1.773	0.699	-1.53	0.37	0.49	1.25
1	1/8	2.029	1.252	-0.62	0.36	0.62	1.01
2	1/2	2.029	1.252	-0.62	0.36	0.62	1.01
3	3/4	2.029	1.252	-0.62	0.57	0.51	0.82
3	1	2.029	1.252	-0.62	0.43	0.59	0.95
2	1/4	2.531	2.259	-0.12	0.28	0.79	0.89
3	1/2	2.531	2.259	-0.12	0.36	0.75	0.84
4	3/4	2.531	2.259	-0.12	0.46	0.69	0.77
4	1	2.531	2.259	-0.12	0.28	0.79	0.89
5	1	2.531	2.259	-0.12	0.54	0.63	0.71
2	1/8	3.280	3.659	0.10	0.24	0.91	0.82
3	1/4	3.280	3.659	0.10	0.33	0.86	0.77
4	1/2	3.280	3.659	0.10	0.24	0.91	0.82
5	3/4	3.280	3.659	0.10	0.27	0.89	0.80
5	1/2	3.444	3.979	0.13	0.45	0.79	0.68
3	1/8	4.222	5.371	0.21	0.36	0.89	0.70
4	1/4	4.222	5.371	0.21	0.29	0.94	0.74
5	1/4	5.226	7.085	0.26	0.25	1.00	0.73
4	1/8	5.910	8.120	0.27	0.25	1.01	0.73
5	1/8	8.496	11.760	0.27	-0.00	1.17	0.84

Numerical Issues

- We used AMPL and NEOS
- Algorithms
 - FilterSQP was most reliable
 - SNOPT was pretty reliable
 - IPOPT stopped early - interior point method is too loose
 - MINOS often failed - unknown reasons
- Global optimization issues
 - Successful algorithms agreed
 - Small deviation examples found qualitatively similar results

Extensions and Conclusions

- Possible heterogeneities
 - $(c - a)^{1-\gamma} / (1 - \gamma) - \psi l^{\eta+1} / (1 + \eta)$ – parameters (a, γ, ψ, η)
 - Age
 - * All people use the same tax tables.
 - * Is low income an indicator of low wage profile or being young and at beginning of wage profile?
 - * Since age is not used, we need to adapt revelation principle
 - Assets holdings
 - Idiosyncratic wage risk
- Real economies have multidimensional populations - need to examine similar models.

Computation of Moral-Hazard Problems

Che-Lin Su* Kenneth L. Judd†

This Draft: June 28, 2005

Abstract

We study computational aspects of moral-hazard problems. In particular, we consider deterministic contracts as well as contracts with action and/or compensation lotteries, and formulate each case as a mathematical program with equilibrium constraints (MPEC). We investigate and compare solution properties of the MPEC approach to that of the linear programming (LP) approach with lotteries. We propose a hybrid procedure that combines the best features of both. The hybrid procedure obtains a solution that is, if not global, at least as good as an LP solution. It also preserves the fast local convergence property by applying the SQP algorithm to MPECs. The numerical results on an example show that the hybrid procedure outperforms the LP approach in both computational time and solution quality in terms of the optimal objective value.

1 Introduction

We study mathematical programming approaches to solve moral-hazard problems. More specifically, we formulate moral-hazard problems with finitely many action choices, including the basic deterministic models and models with lotteries, as *mathematical programs with equilibrium constraints* (MPECs). One advantage of using an MPEC formulation is that the size of resulting program is often orders of magnitude smaller than the linear programs derived from the LP lotteries approach [16, 17]. This feature makes the MPEC approach an appealing alternative when solving a large-scale linear program is computationally infeasible because of limitations on computer memory or computing time.

*Department of Management Science and Engineering, Stanford University (clsu@stanford.edu). The work of this author's research is supported by the Jerome Kaseberg Doolan Fellowship, Stanford University. This author thanks Prof. Jong-Shi Pang for suggesting and Prof. Richard W. Cottle for encouraging this collaboration with Dr. Kenneth L. Judd. This author is grateful to Dr. Sven Leyffer and Dr. Todd Munson for their valuable discussions during his visit to Argonne National Laboratory.

†Hoover Institution, Stanford, CA 94305 (judd@hoover.stanford.edu) and NBER.

The moral-hazard model studies the relationship between a principal (leader) and an agent (follower) in situations in which the principal can neither observe nor verify an agent's action. The model is formulated as a bilevel program, in which the principal's upper-level decision takes the agent's best response to the principal's decision into account. Bilevel programs are generally difficult mathematical problems and much research in the economics literature has been devoted to analyzing and characterizing solutions of the moral-hazard model (see Grossman and Hart [6] and the references therein). When the agent's set of actions is a continuum, an intuitive approach to simplifying the model is to assume the agent's optimal action lies in the interior of the action set. One then can treat the agent's problem as an unconstrained maximization problem and replace it by the first-order optimality conditions. This is called the *first-order approach* in the economics literature. However, Mirrlees [11] showed that the first-order approach may be invalid because the lower-level agent's problem is not necessarily a concave maximization program and that the optimal solution may fail to be unique and interior. Consequently, a sequence of papers [18, 7, 8] has developed conditions under which the first-order approach is valid. Unfortunately, these conditions are often more restrictive than is desirable.

In general, if the lower-level problem in a bilevel program is a convex minimization (or concave maximization) problem, one can replace the lower-level problem by the first-order optimality conditions, which are both necessary and sufficient, and reformulate the original bilevel problem as an MPEC. This idea is similar to the first-order approach to the moral-hazard problem with one notable difference: MPEC formulations include complementarity constraints. The first-order approach assumes that the solution to the agent's problem lies in the interior of the action set, and hence, one can treat the agent's problem as an unconstrained maximization problem. This assumption may also avoid issues associated with the failure of the constraint qualification at a solution. General bilevel programs do not make an interior solution assumption. As a result, the complementarity conditions associated with the Karush-Kuhn-Tucker multipliers for inequality constraints would appear in the first-order optimality conditions for the lower-level program. MPECs also arise in many applications in engineering (e.g., transportation, contact problems, mechanical structure design) and economics (Stackelberg games, optimal taxation problems). One well-known theoretical difficulty with MPECs is that the standard constraint qualifications, such as the linear independence constraint qualification and the Mangasarian-Fromovitz constraint qualification, fail at every feasible point. An extensive literature has developed to refine constraint qualifications and stationarity conditions for MPECs; see Scheel and Scholtes [20] and the references therein. We also refer to the two-volume monograph by Facchinei and Pang [2] for theory and applications of equilibrium problems and to the monographs by Luo et al. [10] and Outrata et al. [14] for more details on MPEC theory and applications.

The failure of the constraint qualification conditions means that the set of Lagrange multipliers is unbounded and that conventional numerical optimization software may fail to converge to a solution. Economists have avoided these numerical problems by reformulating the moral-hazard problem as a linear program involving lotteries over a finite set of outcomes. See Townsend [23, 24] and Prescott [16, 17]. While this approach avoids the constraint

qualification problems, it does so by restricting aspects of the contract, such as consumption, to a finite set of possible choices even though a continuous choice formulation would be economically more natural.

The purpose of this paper is twofold: (1) To introduce to the economics community the MPEC approach, or more generally, advanced equilibrium programming approaches, to the moral hazard problem; (2) To present an interesting and important class of incentive problems in economics to the mathematical programming community. Many incentive problems, such as contract design, optimal taxation and regulation, and multiproduct pricing, can be naturally formulated as an MPEC or an *equilibrium problem with equilibrium constraints* (EPEC) [21, 22]. This greatly extends the applications of equilibrium programming to one of the most active research topics in economics in past three decades. The need for a global solution for these economical problems provides a motivation for optimization community to develop efficient global optimization algorithms for MPECs and EPECs.

The remainder of this paper is organized as follows. In the next section, we describe the basic moral-hazard model and formulate it as a mixed-integer nonlinear program and as an MPEC. In Section 3, we consider moral-hazard problems with action lotteries, with compensation lotteries, and with a combination of both. We derive MPEC formulations for each of these cases. We also compare the properties of the MPEC approach and the LP lottery approach. In Section 5, we develop a hybrid approach that preserves the robustness of global solution from the LP approach and the fast local convergence of the MPEC approach. The numerical efficiency of the hybrid approach in both computational speed and robustness of the solution is illustrated in an example.

2 The Basic Moral-Hazard Model

2.1 The Deterministic Contract

We consider a moral-hazard model in which the agent chooses an action from a finite set $\mathcal{A} = \{a_1, \dots, a_M\}$. The outcome can be one of N alternatives. Let $\mathcal{Q} = \{q_1, \dots, q_N\}$ denote the outcome space where the outcomes are dollar returns to the principal ordered from smallest to largest.

The principal can only observe the outcome, not the agent's action. However, the stochastic relationship between actions and outcomes, which is often called a *production technology*, is common knowledge to both the principal and the agent. Usually, the production technology is exogenously described by the probability distribution function, $p(q|a)$, which presents the probability of outcome $q \in \mathcal{Q}$ occurring given that action a is taken. We assume $p(q|a) > 0$ for all $q \in \mathcal{Q}$ and $a \in \mathcal{A}$; this is called the *full-support assumption*.

Since the agent's action is not observable to the principal, the payment to the agent is

only based on the outcome observed by the principal. Let $\mathcal{C} \subset R$ be the set of all possible compensations.

Definition 1 A *compensation schedule* $c = (c(q_1), \dots, c(q_N)) \in R^N$ is an agreement between the principal and the agent such that $c(q) \in \mathcal{C}$ is the payoff to the agent from the principal if outcome $q \in \mathcal{Q}$ is observed.

The agent's utility $u(x, a)$ is a function of the payment $x \in R$ received from the principal and of his action $a \in \mathcal{A}$. The principal's utility $w(q - x)$ is a function over net income $q - x$ for $q \in \mathcal{Q}$. We let $W(c, a)$ and $U(c, a)$ denote the *expected utility* to the principal and agent, respectively, of a compensation schedule $c \in R^N$ when the agent chooses action $a \in \mathcal{A}$, i.e.,

$$\begin{aligned} W(c, a) &= \sum_{q \in \mathcal{Q}} p(q | a) w(q - c(q)), \\ U(c, a) &= \sum_{q \in \mathcal{Q}} p(q | a) u(c(q), a). \end{aligned} \tag{1}$$

Definition 2 A *deterministic contract* (proposed by the principal) consists of a recommended action $a \in \mathcal{A}$ to the agent and a compensation schedule $c \in R^N$.

The contract has to satisfy two conditions to be accepted by the agent. The first condition is the *participation constraint*. It states that the contract must give the agent an expected utility no less than a required utility level U^* :

$$U(c, a) \geq U^*. \tag{2}$$

The value U^* represents the highest utility the agent can receive from other activities if he does not sign the contract.

Second, the contract must be *incentive compatible* to the agent; it has to provide incentives for the agent not to deviate from the recommended action. In particular, given the compensation schedule c , the recommended action a must be optimal from the agent's perspective and maximizes the agent's expected utility function. The *incentive compatibility constraint* is given as follows:

$$a \in \operatorname{argmax}\{U(c, a) : a \in \mathcal{A}\}. \tag{3}$$

For a given U^* , a *feasible* contract satisfies the participation constraint (2) and the incentive compatibility constraint (3). The objective of the principal is to find an *optimal deterministic contract*, a feasible contract that maximize his expected utility. A mathematical program for finding an optimal deterministic contract (c^*, a^*) is:

$$\begin{aligned} &\operatorname{maximize}_{(c,a)} W(c, a) \\ &\text{subject to} \quad U(c, a) \geq U^*, \\ &\quad a \in \operatorname{argmax}\{U(c, a) : a \in \mathcal{A}\}. \end{aligned} \tag{4}$$

Since there are only finitely many actions in \mathcal{A} , the incentive compatibility constraint (3) can be presented as the following set of inequalities:

$$U(c, a) \geq U(c, a_i), \quad \text{for } i = 1, \dots, M. \quad (5)$$

These constraints ensure that the agent's expected utility obtained from choosing the recommendation action is no worse than that of choosing other actions. Replacing (3) by the set of inequalities (5), we have an equivalent formulation of the optimal contract problem:

$$\begin{aligned} & \text{maximize}_{(c,a)} && W(c, a) \\ & \text{subject to} && U(c, a) \geq U^*, \\ & && U(c, a) \geq U(c, a_1), \\ & && \vdots \\ & && U(c, a) \geq U(c, a_M), \\ & && a \in \mathcal{A} = \{a_1, \dots, a_M\}. \end{aligned} \quad (6)$$

2.2 A Mixed-Integer Nonlinear Programming Formulation

The optimal contract problem (6) can be formulated as a mixed-integer nonlinear program. Associated with each action $a_i \in \mathcal{A}$, we introduce a binary variable $y_i \in \{0, 1\}$. Let $y = (y_1, \dots, y_M) \in R^M$ and let e_M denote the vector of all ones in R^M . To ease the notation, we define

$$\begin{aligned} U(c) &= (U(c, a_1), \dots, U(c, a_M)) \in R^M, \\ W(c) &= (W(c, a_1), \dots, W(c, a_M)) \in R^M. \end{aligned} \quad (7)$$

The mixed-integer nonlinear programming formulation for the optimal contract problem (6) is

$$\begin{aligned} & \text{maximize}_{(c,y)} && y^T W(c) \\ & \text{subject to} && y^T U(c) \geq U^*, \\ & && y^T U(c) \geq U(c, a_1), \\ & && \vdots \\ & && y^T U(c) \geq U(c, a_M), \\ & && e_M^T y = 1, \\ & && y_i \in \{0, 1\} \quad \forall i = 1, \dots, M. \end{aligned} \quad (8)$$

The above problem has N nonlinear variables, M binary variables, one linear constraint and $(M + 1)$ nonlinear constraints. To solve a mixed-integer nonlinear program, one can use MINLP [3], BARON [19] or other solvers developed for this class of programs. For (8), since the agent will choose one and only one action, the number of possible combinations on the binary vector y is only M . One then can solve (8) by solving M nonlinear programs

with $y_i = 1$, and $y_j = 0$ in the i -th nonlinear program, as Grossman and Hart suggested in [6] for the case where the principal is risk averse. They further point out that each nonlinear program is a convex program if the agent's utility function $u(x, a)$ can be written as $G(a) + K(a)V(x)$, where (1) V is real-valued, strictly increasing, concave function defined on some open interval $\mathcal{I} = (\underline{I}, \bar{I}) \subset R$; (2) $\lim_{x \rightarrow \underline{I}} V(x) = -\infty$; (3) G, K are real-valued functions defined on \mathcal{A} and K is strictly positive; (4) $u(x, a) \geq u(x, \hat{a}) \Rightarrow u(\hat{x}, a) \geq u(\hat{x}, \hat{a})$, for all $a, \hat{a} \in \mathcal{A}$, and $x, \hat{x} \in \mathcal{I}$. The above assumption implies that the agent's preferences over income lotteries are independent of his action.

2.3 An MPEC Formulation

In general, a mixed-integer nonlinear program is a difficult optimization problem. Below, by considering a mixed-strategy reformulation of the incentive compatibility constraints for the agent, we can reformulate the optimal contract problem (6) as a mathematical program with equilibrium constraints (MPEC); see [10].

For $i = 1, \dots, M$, let δ_i denote the probability that the agent will choose action a_i . Then, given the compensation schedule c , the agent chooses a mixed strategy profile $\delta^* = (\delta_1^*, \dots, \delta_M^*) \in R^M$ such that

$$\delta^* \in \operatorname{argmax} \left\{ \sum_{k=1}^M \delta_k U(c, a_k) : e_M^T \delta = 1, \delta \geq 0 \right\}. \quad (9)$$

Observe that the agent's mixed-strategy problem (9) is a linear program, and hence, its optimality conditions are necessary and sufficient.

The following lemma states the relationship between the optimal pure strategy a_i and the optimal mixed strategy δ^* .

Lemma 1 Given a compensation schedule $\bar{c} \in R^N$, the agent's action $a_i \in \mathcal{A}$ is optimal to problem (3) iff there exists an optimal mixed strategy profile δ^* to problem (9) such that

$$\begin{aligned} \delta_i^* &> 0, \\ \sum_{k=1}^M \delta_k^* U(\bar{c}, a_k) &= U(\bar{c}, a_i), \\ e_M^T \delta^* &= 1, \quad \delta^* \geq 0. \end{aligned}$$

Proof If a_i is an optimal action of (3), then let $\delta^* = e_i$, where e_i is the i -th column in an identity matrix of order M . It is easy to verify that all the conditions for δ^* are satisfied. Conversely, if a_i is not an optimal solution of (3), then there exists an optimal action a_j such that $U(\bar{c}, a_j) > U(\bar{c}, a_i)$. Let $\tilde{\delta} = e_j$. Then $\tilde{\delta}^T U(c) = U(\bar{c}, a_j) = U(\bar{c}, a_i) > \delta^{*T} U(c)$. We have a contradiction. \square

Let $u \perp v$ indicate orthogonality of vectors u and v , i.e., $u^T v = 0$. An observation following from Lemma 1 is stated below.

Lemma 2 Given a compensation schedule $c \in R^N$, a mixed strategy profile δ is optimal to the linear program (9) iff

$$0 \leq \delta \perp (\delta^T U(c)) e_M - U(c) \geq 0, \quad e_M^T \delta = 1. \quad (10)$$

Proof. This follows from the optimality conditions and the strong duality theorem for the LP (9). \square

Substituting the incentive compatibility constraint (5) by the system (10) and replacing $W(c, a)$ and $U(c, a)$ by $\delta^T W(c)$ and $\delta^T U(c)$, respectively, we derive an MPEC formulation of the principal's problem (6):

$$\begin{aligned} & \text{maximize}_{(c, \delta)} && \delta^T W(c) \\ & \text{subject to} && \delta^T U(c) \geq U^*, \\ & && e_M^T \delta = 1, \\ & && 0 \leq \delta \perp (\delta^T U(c)) e_M - U(c) \geq 0. \end{aligned} \quad (11)$$

To illustrate the failure of constraint qualification at any feasible point of an MPEC, we consider the feasible region $\mathcal{F}_1 = \{(x, y) \in R^2 \mid x \geq 0, y \geq 0, xy = 0\}$. At the point $(\bar{x}, \bar{y}) = (0, 2)$, the first constraint $x \geq 0$ and the third constraint $xy = 0$ are binding. Vectors of the gradient of the binding constraints at (\bar{x}, \bar{y}) are $(1, 0)$ and $(2, 0)$, which are dependent. It is easy to verify that the gradient vectors of the binding constraints are indeed dependent at other feasible points.




Fig. 1. The feasible region $\mathcal{F}_1 = \{(x, y) \mid x \geq 0, y \geq 0, xy = 0\}$.

The following lemma states the relationship between the optimal solutions for the principal-agent problems (6) and the corresponding MPEC formulation (11).

Theorem 1 If (c^*, δ^*) is an optimal solution for the MPEC (11), then (c^*, a_i^*) , where $i \in \{j : \delta_j^* > 0\}$, is an optimal solution for the problem (6). Conversely, if (c^*, a_i^*) is an optimal solution for the problem (6), then (c^*, e_i) is an optimal solution for the MPEC (11).

Proof The statement follows directly from Lemma 2. □

The MPEC (11) has $(N + M)$ variables, 1 linear constraint, 1 nonlinear constraint, and M complementarity constraints. Hence, the size of the problem grows linearly in the number of the outcomes and actions. As we will see in Section 3.4, this feature is the main advantage of using the MPEC approach rather than the LP lotteries approach.

3 Moral-Hazard Problems with Lotteries

In this section, we study moral-hazard problems with lotteries. In particular, we consider action lotteries, compensation lotteries, and a combination of both. For each case, we first give definitions for the associated lotteries and then derive the nonlinear programming or MPEC formulation.

3.1 The Contract with Action Lotteries

Definition 3 A *contract with action lotteries* is a probability distribution over actions, $\pi(a)$, and a compensation schedule $c(a) = (c(q_1, a), \dots, c(q_N, a)) \in R^N$ for all $a \in \mathcal{A}$. The compensation schedule $c(a)$ is an *agreement* between the principal and the agent such that $c(q, a) \in \mathcal{C}$ is the *payoff* to the agent from the principal if outcome $q \in \mathcal{Q}$ is observed and the action $a \in \mathcal{A}$ is recommended by the principal.

In the definition of a contract with action lotteries, the compensation schedule $c(a)$ is contingent on both the outcome and the agent's action. Given this definition, one might raise the following question: if the principal can only observe the outcome, not the agent's action, is it reasonable to have the compensation schedule $c(a)$ contingent on the action chosen by the agent? After all, the principal does not know which action is implemented by the agent. One economic justification is as follows. Suppose that the principal and the agent sign a total of M contracts, each with different recommended action $a \in \mathcal{A}$ and compensation schedule $c(a)$ as a function of the recommended action, a . Then, the principal and the agent would go to an authority or a third party to conduct a lottery with probability distribution function $\pi(a)$ on which contract would be implemented on that day. If the i -th contract is drawn from the lottery, then the third party would inform both the principal and the agent that the recommended action for that day is a_i with the compensation schedule $c(a_i)$.

Arnott and Stiglitz [1] use *ex ante randomization* for action lotteries. This terminology refers to the situation that a random contract occurs before the recommended action is chosen. They demonstrate that the action lotteries will result in a welfare improvement if the principal's expected utility is nonconcave in the agent's expected utility. However, it is not clear which the sufficient conditions would be needed for the statement in the assumption to be true.

$(\pi^*(a), c^*(a))_{a \in \mathcal{A}}$ is then a solution to the following nonlinear program:

$$\begin{aligned}
& \text{maximize} && \sum_{a \in \mathcal{A}} \pi(a) W(c(a), a) \\
& \text{subject to} && \sum_{a \in \mathcal{A}} \pi(a) U(c(a), a) \geq U^*, \\
& && \sum_{a \in \mathcal{A}} \pi(a) = 1, \\
& && \pi(a) (U(c(a), a) - U(c(a), \tilde{a})) \geq 0, \quad \forall (a, \tilde{a} (\neq a)) \in \mathcal{A} \times \mathcal{A}, \\
& && \pi(a) \geq 0, \quad \forall a \in \mathcal{A}.
\end{aligned} \tag{18}$$

The nonlinear program (18) has $(N * M + M)$ variables and $(M * (M - 1) + 2)$ constraints. In addition, its feasible region is highly nonconvex because of the last two sets of constraints in (18). As shown in the following graph, the feasible region $\mathcal{F}_2 = \{(x, y) \mid xy \geq 0, x \geq 0\}$ is the union of the first quadrant and the y -axis. Furthermore, the standard nonlinear programming constraint qualification fails to hold at every point on the y -axis.




Fig. 2. The feasible region $\mathcal{F}_2 = \{(x, y) \mid xy \geq 0, x \geq 0\}$.

MPEC formulations

Below, we introduce an MPEC formulation for the star-shaped problem (18). We first show that constraints of a star-shaped set $\mathcal{Z}_1 = \{z \in R^n \mid g_i(z) \geq 0, g_i(z) h_i(z) \geq 0, i = 1, \dots, m\}$ can be rewritten as complementarity constraints by introducing additional variables.

Proposition 1 A point z is in $\mathcal{Z}_1 = \{z \in R^n \mid g_i(z) \geq 0, g_i(z) h_i(z) \geq 0, i = 1, \dots, m\}$ iff there exists an s such that (z, s) is in $\mathcal{Z}_2 = \{(z, s) \in R^{n+m} \mid 0 \leq g(z) \perp s \geq 0, h(z) \geq -s\}$.

Proof. Suppose that z is in \mathcal{Z}_1 . If $g_i(z) > 0$, choose $s_i = 0$; if $g_i(z) = 0$, choose $s_i = -h_i(z)$. Then (z, s) is in \mathcal{Z}_2 . Conversely, if (z, s) is in \mathcal{Z}_2 , then $g_i(z) h_i(z) \geq g_i(z) (-s_i) = 0$ for all $i = 1, \dots, m$. Hence, the point z is in \mathcal{Z}_1 . \square

Following Proposition 1, we introduce a variable $s(a)$ for each $a \in \mathcal{A}$ for the incentive compatibility constraints in (18). We then obtain the following MPEC formulation with variables $(\pi(a), c(a), s(a))_{a \in \mathcal{A}}$ for the optimal contract with action lottery problem:

$$\begin{aligned}
& \text{maximize} && \sum_{a \in \mathcal{A}} \pi(a) W(c(a), a) \\
& \text{subject to} && \sum_{a \in \mathcal{A}} \pi(a) U(c(a), a) \geq U^*, \\
& && \sum_{a \in \mathcal{A}} \pi(a) = 1, \\
& && U(c(a), a) - U(c(a), \tilde{a}) + s(a) \geq 0, \quad \forall (a, \tilde{a} (\neq a)) \in \mathcal{A} \times \mathcal{A}, \\
& && 0 \leq \pi(a) \perp s(a) \geq 0.
\end{aligned} \tag{19}$$

Allowing the compensation schedules to be dependent on the agent's action will increase the principal's expected utility; see Theorem 2 below. The difference between the optimal objective value of the NLP (18) (or the MPEC(19)) and that of the MPEC (11) characterizes the principal's improved welfare from using an optimal contract action lotteries.

Theorem 2 The principal prefers an optimal contract with action lotteries to an optimal deterministic contract. His expected utility from choosing an optimal contract with action lotteries will be at least as good as that from choosing an optimal deterministic contract.

Proof This is clear. □

3.2 The Contract with Compensation Lotteries

Definition 4 For any outcome $q \in \mathcal{Q}$, a *randomized compensation* $\tilde{c}(q)$ is a random variable on the set of compensations \mathcal{C} with a probability measure $F(\cdot)$.

Remark If the set of compensations \mathcal{C} is a closed interval $[\underline{c}, \bar{c}] \in \mathbb{R}$, then the measure of $\tilde{c}(q)$ is a cumulative density function (cdf) $F : [\underline{c}, \bar{c}] \rightarrow [0, 1]$ with $F(\underline{c}) = 0$ and $F(\bar{c}) = 1$. In addition, $F(\cdot)$ is nondecreasing and right-continuous.

To simplify the analysis, we assume that every randomized compensation $\tilde{c}(q)$ has finite support.

Assumption 1: Finite support for randomized compensation. For all $q \in \mathcal{Q}$, the randomized compensation $\tilde{c}(q)$ has finite support over an unknown set $\{c_1(q), c_2(q), \dots, c_L(q)\}$ with a known L .

An immediate consequence of Assumption 1 is that we can write $\tilde{c}(q) = c_i(q)$ with probability $p_i(q) > 0$ for all $i = 1, \dots, L$ and $q \in \mathcal{Q}$. In addition, we have $\sum_{i=1}^L p_i(q) = 1$ for

all $q \in \mathcal{Q}$. Notice that both $(c_i(q))_{i=1}^L \in R^L$ and $(p_i(q))_{i=1}^L \in R^L$ are endogenous variables and will be chosen by the principal.

Definition 5 A *compensation lottery* is a randomized compensation schedule $\tilde{c} = (\tilde{c}(q_1), \dots, \tilde{c}(q_N)) \in R^N$, in which $\tilde{c}(q)$ is a randomized compensation satisfying Assumption 1 for all $q \in \mathcal{Q}$.

Definition 6 A *contract with compensation lotteries* consists of a recommended action a to the agent and a randomized compensation schedule $\tilde{c} = (\tilde{c}(q_1), \dots, \tilde{c}(q_N)) \in R^N$.

Let $c^q = (c_i(q))_{i=1}^L \in R^L$ and $p^q = (p_i(q))_{i=1}^L \in R^L$. Given that the outcome q is observed by the principal, we let $\mathbf{w}(c^q, p^q)$ denote the principal's expected utility with respect to a randomized compensation $\tilde{c}(q)$, i.e.,

$$\mathbf{w}(c^q, p^q) = \mathbb{E} w(q - \tilde{c}(q)) = \sum_{i=1}^L p_i(q) w(q - c_i(q)).$$

With a randomized compensation schedule \tilde{c} and a recommended action a , the principal's expected utility then becomes

$$\mathbb{E} W(\tilde{c}, a) = \sum_{q \in \mathcal{Q}} p(q | a) \left(\sum_{i=1}^L p_i(q) w(q - c_i(q)) \right) = \sum_{q \in \mathcal{Q}} p(q | a) \mathbf{w}(c^q, p^q). \quad (20)$$

Similarly, given a recommended action a , we let $\mathbf{u}(c^q, p^q, a)$ denote the agent's expected utility with respect to $\tilde{c}(q)$ for the observed outcome q :

$$\mathbf{u}(c^q, p^q, a) = \mathbb{E} u(\tilde{c}(q), a) = \sum_{i=1}^L p_i(q) u(c_i(q), a).$$

The agent's expected utility with a randomized compensation schedule \tilde{c} and a recommended action a is

$$\mathbb{E} U(\tilde{c}, a) = \sum_{q \in \mathcal{Q}} p(q | a) \left(\sum_{i=1}^L p_i(q) u(c_i(q), a) \right) = \sum_{q \in \mathcal{Q}} p(q | a) \mathbf{u}(c^q, p^q, a). \quad (21)$$

To further simplify notation, we use $c_{\mathcal{Q}} = (c^q)_{q \in \mathcal{Q}}$ and $p_{\mathcal{Q}} = (p^q)_{q \in \mathcal{Q}}$ to denote the collection of variables c^q and p^q , respectively. We also let $\mathbf{W}(c_{\mathcal{Q}}, p_{\mathcal{Q}}, a)$ denote the principal's expected utility $\mathbb{E} W(\tilde{c}, a)$ as defined in (20), and similarly, $\mathbf{U}(c_{\mathcal{Q}}, p_{\mathcal{Q}}, a)$ for $\mathbb{E} U(\tilde{c}, a)$ as in (21).

An optimal contract with compensation lotteries $(c_{\mathcal{Q}}^*, p_{\mathcal{Q}}^*, a^*)$ is a solution to the following

problem:

$$\begin{aligned}
& \text{maximize} && \mathbf{W}(c_Q, p_Q, a) \\
& \text{subject to} && \mathbf{U}(c_Q, p_Q, a) \geq U^*, \\
& && \mathbf{U}(c_Q, p_Q, a) \geq \mathbf{U}(c_Q, p_Q, a_1), \\
& && \vdots \\
& && \mathbf{U}(c_Q, p_Q, a) \geq \mathbf{U}(c_Q, p_Q, a_M), \\
& && a \in \mathcal{A} = \{a_1, \dots, a_M\}.
\end{aligned} \tag{22}$$

Define

$$\begin{aligned}
\mathbf{W}(c_Q, p_Q) &= (\mathbf{W}(c_Q, p_Q, a_1), \dots, \mathbf{W}(c_Q, p_Q, a_M)) \in R^M, \\
\mathbf{U}(c_Q, p_Q) &= (\mathbf{U}(c_Q, p_Q, a_1), \dots, \mathbf{U}(c_Q, p_Q, a_M)) \in R^M.
\end{aligned}$$

Following the derivation as in Section 2, we can reformulate the program for an optimal contract with compensation lotteries (22) as a mixed-integer nonlinear program with decision variables (c_Q, p_Q) and $y = (y_i)_{i=1}^M$:

$$\begin{aligned}
& \text{maximize} && y^T \mathbf{W}(c_Q, p_Q) \\
& \text{subject to} && y^T \mathbf{U}(c_Q, p_Q) \geq U^*, \\
& && y^T \mathbf{U}(c_Q, p_Q) \geq \mathbf{U}(c_Q, p_Q, a_1), \\
& && \vdots \\
& && y^T \mathbf{U}(c_Q, p_Q) \geq \mathbf{U}(c_Q, p_Q, a_M), \\
& && e_M^T y = 1, \\
& && y_i \in \{0, 1\} \quad \forall i = 1, \dots, M,
\end{aligned} \tag{23}$$

Similarly, the MPEC formulation with decision variables (c_Q, p_Q) and $\delta \in R^M$ is

$$\begin{aligned}
& \text{maximize} && \delta^T \mathbf{W}(c_Q, p_Q) \\
& \text{subject to} && \delta^T \mathbf{U}(c_Q, p_Q) \geq U^*, \\
& && e_M^T \delta = 1, \\
& && 0 \leq \delta \perp \left(\delta^T \mathbf{U}(c_Q, p_Q) \right) e_M - \mathbf{U}(c_Q, p_Q) \geq 0.
\end{aligned} \tag{24}$$

Arnott and Stiglitz [1] call the compensation lotteries *ex post randomization*; this refers to the situation where the random compensation occurs after the recommended action is chosen or implemented. They show that if the agent is risk averse and his utility function is separable, and if the principal is risk neutral, then the compensation lotteries are not desirable.

3.3 The Contract with Action and Compensation Lotteries

Definition 7 A contract with action and compensation lotteries is a probability distribution over actions, $\pi(a)$, and a randomized compensation schedule $\tilde{c}(a) = (\tilde{c}(q_1, a), \dots, \tilde{c}(q_N, a)) \in R^N$ for every $a \in \mathcal{A}$. The randomized compensation schedule $c(a)$ is an agreement between the principal and the agent such that $\tilde{c}(q, a) \in \mathcal{C}$ is a randomized compensation to the agent from the principal if outcome $q \in \mathcal{Q}$ is observed and the action $a \in \mathcal{A}$ is recommended by the principal.

Assumption 2 For every action $a \in \mathcal{A}$, the randomized compensation schedule $\tilde{c}(q, a)$ satisfies the finite support assumption (Assumption 1) for all $q \in \mathcal{Q}$.

With Assumption 2, the notation $c^q(a), p^q(a), c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a)$ is analogous to what we have defined in Section 3.1 and 3.2. Without repeating the same derivation process described earlier, we give the star-shaped formulation with variables $(\pi(a), c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a))_{a \in \mathcal{A}}$ for the optimal contract with action and compensation lotteries problem:

$$\begin{aligned}
& \text{maximize} && \sum_{a \in \mathcal{A}} \pi(a) \mathbf{W}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), a) \\
& \text{subject to} && \sum_{a \in \mathcal{A}} \pi(a) \mathbf{U}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), a) \geq U^*, \\
& && \sum_{a \in \mathcal{A}} \pi(a) = 1, \\
& && \pi(a) (\mathbf{U}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), a) - \mathbf{U}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), \tilde{a})) \geq 0, \quad \forall (a, \tilde{a}) \in \mathcal{A} \times \mathcal{A}, \\
& && \pi(a) \geq 0.
\end{aligned} \tag{25}$$

Following the derivation in Section 3.1, an equivalent MPEC formulation is with variables $(\pi(a), c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), s(a))_{a \in \mathcal{A}}$:

$$\begin{aligned}
& \text{maximize} && \sum_{a \in \mathcal{A}} \pi(a) \mathbf{W}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), a) \\
& \text{subject to} && \sum_{a \in \mathcal{A}} \pi(a) \mathbf{U}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), a) \geq U^*, \\
& && \sum_{a \in \mathcal{A}} \pi(a) = 1, \\
& && \mathbf{U}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), a) - \mathbf{U}(c_{\mathcal{Q}}(a), p_{\mathcal{Q}}(a), \tilde{a}) + s(a) \geq 0, \quad \forall (a, \tilde{a}) \in \mathcal{A} \times \mathcal{A}, \\
& && 0 \leq \pi(a) \perp s(a) \geq 0.
\end{aligned} \tag{26}$$

Linear Programming Approximation

Townsend [23, 24] was among the first to use linear programming techniques to compute static incentive constrained problems. Prescott [16, 17] further apply linear programming specifically to solve moral-hazard problems. A solution obtained by the linear programming

approach is an approximation to a solution to the MPEC (26). Instead of treating $c_{\mathcal{Q}}(a)$ as unknown variables, one can construct a grid Ξ with element ξ to approximate the set \mathcal{C} of compensations. By introducing probability measures associated with the action lotteries on \mathcal{A} and compensation lotteries on Ξ , one can then approximate a solution to the moral-hazard problem with lotteries (26) by solving a linear program. More specifically, the principal chooses probability distributions $\pi(a)$, and $\pi(\xi|q, a)$ over the set of actions \mathcal{A} , the set of outcomes \mathcal{Q} , and the compensation grid Ξ . One then can reformulate the resulting nonlinear program into a linear program with decision variables $\pi = (\pi(\xi, q, a))_{\xi \in \Xi, q \in \mathcal{Q}, a \in \mathcal{A}}$:

$$\begin{aligned}
& \text{maximize}_{(\pi)} \quad \sum_{\xi, q, a} w(q - \xi) \pi(\xi, q, a) \\
& \text{subject to} \quad \sum_{\xi, q, a} u(\xi, a) \pi(\xi, q, a) \geq U^*, \\
& \quad \sum_{\xi, q} u(\xi, a) \pi(\xi, q, a) \geq \sum_{\xi, q} u(\xi, \tilde{a}) \frac{p(q|\tilde{a})}{p(q|a)} \pi(\xi, q, a) \quad \forall (a, \tilde{a} (\neq a)) \in \mathcal{A} \times \mathcal{A}, \\
& \quad \sum_{\xi} \pi(\xi, \tilde{q}, \tilde{a}) = p(\tilde{q}|\tilde{a}) \sum_{\xi, q} \pi(\xi, q, \tilde{a}) \quad \forall (\tilde{q}, \tilde{a}) \in \mathcal{Q} \times \mathcal{A}, \\
& \quad \sum_{\xi, q, a} \pi(\xi, q, a) = 1, \\
& \quad \pi(\xi, q, a) \geq 0 \quad \forall (\xi, q, a) \in \Xi \times \mathcal{Q} \times \mathcal{A}.
\end{aligned} \tag{27}$$

Note that the above linear program has $(|\Xi| * N * M)$ variables and $(M * (N + M - 1) + 2)$ constraints. The size of the linear program will grow enormously and quickly when one chooses a fine grid. For example, if there are 50 actions, 40 outputs, and 500 compensations, then the linear program has one million variables and 4452 constraints. It will become computationally infeasible because of the limitation on computer memory, if not the time required to solve a large-scale linear program. On the other hand, a solution of the LP obtained from a coarse grid will not be satisfactory if an accurate solution is needed. Prescott [17] points out that the constraint matrix of the linear program (27) has block angular structure. As a consequence, one can apply the Dantzig-Wolfe decomposition algorithm to the linear program (27) to reduce the computer memory and computational time. Recall that the MPEC (11) for the optimal contract problem has only $(N + M)$ variables and M complementarity constraints with one linear constraint and one nonlinear constraint. Even with the use of the Dantzig-Wolfe decomposition algorithm to solve LP (27), choosing the “right” grid is still an issue. With the advances in both theory and numerical methods for solving MPECs in the last decade, we believe that the MPEC approach has greater advantages in solving a much smaller problem and in obtaining a more accurate solution.

The error from discretizing set of compensations \mathcal{C} is characterized by the difference between the optimal objective value of LP (27) and that of MPEC (26).

Theorem 3 Assume the agent is risk averse over the payoff received from the principal and his efforts and the principal is risk averse or risk neutral over his net income. Then the optimal objective value of MPEC (26) is at least as good as that of LP (27).

Proof See the proof for Theorem 5.3 in [22]. □

4 A Hybrid Approach toward Global Optimization

One reason that nonconvex programs are not popular among economists is the issue of the need for global solutions. While local search algorithms for solving nonconvex programs have fast convergence properties near a solution, they are designed to find a local solution. Algorithms for solving MPECs are no exception. One heuristic in practice is to solve the same problem with several different starting points. It then becomes a trade-off between the computation time and the quality of the “best” solution found.

Linear programming does not suffer from the global solution issue. However, to obtain an accurate solution to a moral-hazard problem via the linear programming approach, one needs to use a very fine compensation grid. This often leads to large-scale linear programs with millions of variables and tens or hundreds thousands of constraints, which might require excessive computer memory or time.

Certainly, there is a need to develop a global optimization method with fast local convergence for MPECs. Below, we propose a hybrid approach combining both MPECs and linear programming approaches to find a global solution (or at least better than the LP solution) of an optimal contract problem. The motivation for this hybrid method comes from the observation that the optimal objective value of the LP approach from a coarse grid could provide a lower bound on the optimal objective value of the MPEC as well as a good guess on the final recommended action a^* . We can then use this information to exclude some undesired local minimizers and to provide a good starting point when we solve the MPEC (11). This heuristic procedure toward a global solution of the MPEC (11) leads to the following algorithm.

A hybrid method for the optimal contract problem as an MPEC (11)

Step 0: Construct a coarse grid.

Step 1: Solve the LP (27) for the given grid.

$$\text{Step 2: } \left\{ \begin{array}{l} (2.1) : \text{ Compute } \pi(\cdot, \cdot, a) = \sum_{\xi \in \Xi} \sum_{q \in \mathcal{Q}} \pi(\xi, q, a), \quad \forall a \in \mathcal{A}; \\ (2.2) : \text{ Compute } \mathbb{E}[\xi(q)] = \sum_{\xi \in \Xi} \xi \pi(\xi, q, a), \quad \forall q \in \mathcal{Q}; \\ (2.3) : \text{ Set the initial point } c^0 = (\mathbb{E}[\xi(q)])_{q \in \mathcal{Q}} \text{ and } \delta^0 = (\pi(\cdot, \cdot, a))_{a \in \mathcal{A}}; \\ (2.4) : \text{ Solve the MPEC (11) with starting point } (c^0, \delta^0). \end{array} \right.$$

Step 3: Refine the grid and repeat Step 1 and Step 2.

Remark If the starting point from an LP solution is close to the optimal solution of the MPEC (11), then the sequence of iterates generated by the SQP algorithm converges Q-quadratically to the optimal solution. See Proposition 5.2 in Fletcher et al. [4].

One can also develop similar procedures to find global solutions for optimal contract problems with action and/or compensation lotteries. However, the MPECs for contracts with lotteries are much more numerically challenging problems than the MPEC (11) for deterministic contracts.

5 An Example and Numerical Results

To illustrate the use of the mixed-integer nonlinear program (8), the MPEC (11) and the hybrid approaches, and to understand the effect on discretizing the set of compensations \mathcal{C} , we only consider to problems of deterministic contracts without lotteries. We consider a two-outcome example in Karaivanov [9]. Before starting the computational work, we summarize in Table 5.1 the problem characteristics of various approaches to computing the optimal deterministic contracts.

Table 5.1. Problem characteristics of various approaches

	MINLP (8)	MPEC (11)	LP (27)
# of Variables	N	$N + M$	$ \Xi * N * M$
# of Binary Variables	M	–	–
# of Constraints	$M + 2$	2	$M * (N + M - 1) + 2$
# of Complem. Constraints	–	M	–

Example 1: No Action and Compensation Lotteries

Assume the principal is risk neutral with utility $w(q - c(q)) = q - c(q)$, and the agent is risk averse with utility $u(c(q), a) = \frac{c^{1-\gamma}}{1-\gamma} + \kappa \frac{(1-a)^{1-\delta}}{1-\delta}$. Suppose there are only two possible outcomes, e.g., a coin-flip. If the desirable outcome (high sale quantities or high production quantities) happens, then the principal receives $q_H = \$3$; otherwise, he receives $q_L = \$1$. For simplicity, we assume that the set of actions \mathcal{A} consists of M equally-spaced effort levels within the closed interval $[0.01, 0.99]$. The production technology for the high outcome is described by $p(q = q_H|a) = a^\alpha$ with $0 < \alpha < 1$. Note that since 0 and 1 are excluded from the action set \mathcal{A} , the full-support assumption on production technology is satisfied.

The parameter values for the particular instance we solve are given in Table 5.2.

Table 5.2. The value of parameters used in the example.

γ	κ	δ	α	U^*	M
0.5	1	0.5	0.7	1	10

We solve this problem first as a mixed-integer nonlinear program (8) and then as an MPEC (11). For the LP lotteries approach, we start with 20 grid points in the compensation grid (we evenly discretize the compensation set \mathcal{C} into 19 segments) and then increase the size of the compensation grid to 50, 100, 200, \dots , 5000.

We submitted the corresponding AMPL programs to the NEOS server [13]. The mixed-integer nonlinear programs were solved using the MINLP solver [3] on the computer host `newton.mcs.anl.gov`. To obtain fair comparisons between the LP, MPEC, and hybrid approaches, we chose SNOPT [5] to solve the associated mathematical programs. The AMPL programs were solved on the computer host `tate.iems.northwestern.edu`.

Table 5.3 gives the solutions returned by the MINLP solver to the mixed-integer nonlinear program (8). We use $y = 0$ and $y = e_M$ as starting points. In both cases, the MINLP solver returns a solution very quickly. However, it does not guarantee to find a global solution.

Table 5.3. Solutions of the MINLP approach.

Starting Point	# of Regular Variables	# of Binary Variables	# of . Constr.	Solve Time (in sec.)	Objective Value
$y = 0$	2	10	12	0.01	1.864854251
$y = e_M$	2	10	12	0.00	1.877265189

For solving the MPEC (11), we try two different starting points to illustrate the possibility of finding only a local solution. The MPEC solutions are given in Table 5.4 below.

Table 5.4. Solutions of the MPEC approach with two different starting points.

Starting Point	# of Variables	# of Complem. Constraints	Read Time (in sec.)	Solve Time (in sec.)	# of Major Iterations	Objective Value
$\delta = 0$	22	10	0	0.07	45	1.079621424
$\delta = e_M$	22	10	0	0.18	126	1.421561553

The solutions for the LP lottery approach with different compensation grids are given in Table 5.5. Notice that the solve time increases faster than the size of the grid when $|\Xi|$ is of orders 10^5 and higher, while the number of major iterations only increases about 3 times when we increase the grid size 250 times (from $|\Xi| = 20$ to $|\Xi| = 5000$).

Table 5.5. Solutions of the LP approach with 8 different grids (# of constraints = 112).

$ \Xi $	# of Variables	Read Time (in sec.)	Solve Time (in sec.)	# of Iterations	Objective Value
20	400	0.01	0.03	31	1.876085819
50	1000	0.02	0.06	46	1.877252488
100	2000	0.04	0.15	53	1.877252488
200	4000	0.08	0.31	62	1.877254211
500	10000	0.21	0.73	68	1.877263962
1000	20000	0.40	2.14	81	1.877262184
2000	40000	0.83	3.53	71	1.877260460
5000	100000	2.19	11.87	101	1.877262793

Finally, for the hybrid approach, we first use the LP solution from a compensation grid with $|\Xi| = 20$ to construct a starting point for the MPEC (11). As one can see in Table 5.6, with a good starting point, it takes SNOPT only 0.01 seconds to find a solution to the example formulated as the MPEC (11). Furthermore, the optimal objective value is higher than that of the LP solution from a fine compensation grid with $|\Xi| = 5000$.

Table 5.6. Solution of the hybrid approach.

LP $ \Xi $	Read Time (in sec.)	Solve Time (in sec.)	# of Iterations	Objective Value
20	0.01	0.03	31	1.876085819
MPEC Starting Point	Read Time (in sec.)	Solve Time (in sec.)	# of Major Iterations	Objective Value
$\delta_6 = 1, \delta_{i(\neq 6)} = 0$	0.02	0.01	13	1.877265298

6 Conclusions and Future Work

The purpose of this paper is to introduce the MPEC approach and apply it to moral-hazard problems. We have presented MPEC formulations for optimal deterministic contract problems and optimal contract problems with action and/or compensation lotteries. We also formulated the former problem as a mixed-integer nonlinear program. To obtain a global solution, we have proposed a hybrid procedure that combines the LP lottery and the MPEC approaches. In this procedure, the LP solution from a coarse compensation grid provides a good starting point for the MPEC. We can then apply specialized MPEC algorithms with fast local convergence rate to obtain a solution. In a numerical example, we have demonstrated that the hybrid method is more efficient than using only the LP lottery approach, which requires the solution of a sequence of large-scale linear programs. Although we cannot prove that the hybrid approach will guarantee to find a global solution, it finds one better than the solution from the LP lottery approach. We plan to test the numerical performance of

the hybrid procedure on other examples such as the bank regulation example in [16] and the two-dimensional action choice example in [17].

One can extend the MPEC approach to single-principal multiple-agent problems without any difficulty. For the model multiple-principal multiple-agent model [12], it can be formulated as an equilibrium problem with equilibrium constraint (EPEC) [21, 22]. We will investigate these two topics in our future research.

Another important topic we plan to explore is the dynamic moral-hazard problem; see Phelan and Townsend [15]. In the literature, dynamic programming is applied to solve this model. We believe that there is an equivalent nonlinear programming formulation. Analogous to the hybrid procedure proposed in Section 4, an efficient method to solve this dynamic model is to combine dynamic programming and nonlinear programming.

References

- [1] Arnott, A., Stiglitz, J. E. (1998): Randomization with asymmetric information, *The Rand Journal of Economics*, **19**, 344-362.
- [2] Facchinei, F., Pang, J.-S. (2003): *Finite-Dimensional Variational Inequalities and Complementarity Problems*, Springer Verlag, New York.
- [3] Fletcher, R., Leyffer, S. (1999): User manual for MINLP_BB, Department of Mathematics, University of Dundee, UK.
- [4] Fletcher, R., Leyffer, S., Ralph, D., Scholtes, S. (2002): Local convergence of SQP methods for mathematical programs with equilibrium constraints. Numerical Analysis Report NA/209, Department of Mathematics, University of Dundee, UK.
- [5] Gill, P.E., Murray, W., Saunders, M.A. (2002): SNOPT: An SQP algorithm for large-scale constrained optimization, *SIAM Journal on Optimization*, **12**, 979-1006.
- [6] Grossman, S.J., Hart, O.D (1983): An analysis of the principal-agent problem, *Econometrica*, **51**, 7-46.
- [7] Hart, O.D, Holmström, B. (1987): The theory of contracts, in (Bewley, T. eds.) *Advances in Economic Theory: Fifth World Congress*, 7-46, Cambridge University Press, Cambridge, UK.
- [8] Jewitt, I. (1988): Justifying the first-order approach to principal-agent problems, *Econometrica*, **56**, 1177-1190.
- [9] Karaivanov, A. K. (2001): Computing moral hazard programs with lotteries Using Matlab, Working paper, Department of Economics, University of Chicago.

- [10] Luo, Z.-Q., Pang, J.-S., Ralph, D. (1996): *Mathematical Programs with Equilibrium Constraints*, Cambridge University Press, Cambridge, UK.
- [11] Mirrlees, J. A. (1975): The theory of moral-hazard and unobservable behavior: part I, Mimeo, Nuffield College, Oxford University, Oxford, UK.
- [12] Myerson, R. B. (1982): Optimal coordination mechanisms in generalized principal-agent problems, *Journal of Mathematical Economics*, **10**, 67–81.
- [13] The NEOS server for optimization, Webpage: www-neos.mcs.anl.gov/neos/
- [14] Outrata, J., Kocvara, M., Zowe, J. (1998): *Nonsmooth Approach to Optimization Problems with Equilibrium Constraints: Theory, Applications, and Numerical Results*, Kluwer Academic Publishers, Dordrecht, The Netherlands.
- [15] Phelan, C., Townsend, R. M. (1991): Computing multi-period, information-constrained optima, *The Review of Economic Studies*, **58**, 853–881.
- [16] Prescott, E. S. (1999): A primer on moral-hazard models, *Federal Reserve Bank of Richmond Quarterly Review*, **85**, 47–77.
- [17] Prescott, E. S. (2004): Computing solutions to moral-hazard programs using the Dantzig–Wolfe decomposition algorithm, *Journal of Economic Dynamics and Control*, **28**, 777–800.
- [18] Rogerson, W. P. (1985): The first-order approach to principal-agent problems, *Econometrica*, **53**, 1357–1368.
- [19] Sahinidis, N. V., Tawarmalani, M. (2004): BARON 7.2: Global Optimization of Mixed-Integer Nonlinear Programs, *User’s Manual*, Available at <http://www.gams.com/dd/docs/solvers/baron.pdf>.
- [20] Scheel, H., Scholtes, S. (2000): Mathematical programs with equilibrium constraints: stationarity, optimality, and sensitivity, *Mathematics of Operations Research*, **25**, 1–22.
- [21] Su, C.-L. (2004): A sequential NCP algorithm for solving equilibrium problems with equilibrium constraints, Working paper, Department of Management Science and Engineering, Stanford University.
- [22] Su, C.-L. (2005): *Equilibrium Problems with Equilibrium Constraints*, Ph.D. Thesis, Department of Management Science and Engineering, Stanford University, Stanford, CA.
- [23] Townsend, R. M. (1987): Economic organization with limited communication, *American Economic Review*, **77**, 954–971.
- [24] Townsend, R. M. (1988): Information constrained insurance: the revelation principle extended, *Journal of Monetary Economics*, **21**, 411–450.